COVID-19 Update 669 Days Since EOC Activation

January 11, 2022



Case Summary

Table 1. Summary of Cases, Deaths, and Outbreaks by Date Reported

	12/26/2021-1/1/2022	12/19 – 12/25/2021	Cumulative	
Cases	29,536 🔶	12,170	441,885	
Avg Daily Cases	4,219 🔶	1,739	664	
Deaths	27 🖊	36	4,469	
Outbreaks	64 🔶	62	3,273	

Daily Cases Reported

COVID-19 Cases by Date Reported








Testing Positivity %

COVID-19 Laboratory Test Positivity Rate – Previous 12 Months




Includes PCR tests performed by COSD Public Health, hospital, and reference laboratories and reported via Electronic Laboratory Reporting (ELR) and line lists. Excludes invalid, indeterminate, and unsatisfactory results.

Testing Positivity %

COVID-19 Percentage Positive


🍪 🎦 LIVE WELL

Percentage of Positive COVID-19 Cases Among Tests by Date Reported Total Number of Cases = 465,607





COVID-19 Confirmed Hospitalized and ICU Patients 30 Days (12/6 – 1/4)



Invest and Protect Our Community

- Testing available in Chula Vista Monday-Saturday
 - 276 Fourth Ave. (M-F)
 - 1301 Oleander Ave. (Tu-Sat)
- Partnered with Communities Fighting COVID
 - Rapid testing offered several time per month at Orange Library





Testing

December 13, 2021 – January 10, 2022

Test	Administered	
PCR Test (Helix Laboratory)	7,355	
PCR Test (County Laboratory)	663	
Rapid Antigen Test	2,000	
Total	10,018	

Isolation and Quarantine Guidance

COVID-19 Status	Exclusion Period	Return On	Precautions
Tested positive OR Symptomatic (regardless of vaccination status)	5-Day Isolation	Day 6 or later with resolution of fever for at least 24 hours (unmedicated) with a negative rapid antigen test taken day 5 or later and with overall symptom improvement.	Wear a mask for 10 full days around others inside your home or in public.
Close Contact Exposure (unvaccinated or vaccinated booster eligible but not yet boosted)	5-Day Quarantine	Day 6 or later while asymptomatic with a negative test taken day 5 or later.	Wear a mask for 10 full days around others inside your home or in public. Watch for symptoms until 10 days after your exposure; immediately isolate upon any symptom development.
Close Contact Exposure (boosted, vaccinated but not yet booster eligible, or recent infection within 90-days)	Not Applicable (May remain at work)	N/A **Must test on day 5**	Wear a mask for 10 full days around others inside your home or in public. Watch for symptoms until 10 days after your exposure; immediately isolate upon any symptom development.

Invest and Protect Our Community

- Vaccines available at six locations throughout the Chula Vista
 - City Hall, 276 Fourth Ave.
 - Sharp Chula Vista, 752 Medical Center Ct
 - South Bay Pharmacy, 750 Medical Center Court
 - HHSA South Region Public Health Center, 690 Oxford St.
 - Sav-On Pharmacy, 720 3rd Ave
 - COVID Clinic West, 644 Naples St
- Homebound vaccinations ongoing
- Assistance with scheduling appointment



COVID-19 Vaccinations

City of Chula Vista Employees

As of January 6, 2021

Department	% Vaccinated
Administration	100%
Mayor & Council	100%
City Clerk	100%
Library	100%
Economic Development	95%
Parks and Recreation	95%
Information Technology Services	93%
Finance	90%
Engineering	88%
Development Services	88%
Human Resources	87%
City Attorney's Office	87%
Public Works	87%
Animal Care	86%
CBAG	85%
Fire	85%
Police	80%
TOTAL	87%

Use this chart to help determine when to give COVID-19 vaccine based on age or health conditions.

	Dose Age	: 5-11	12-17	18+
Pfizer– Pediatric (5-11)	Primary 2 doses (21 days apart)	1		
	Additional (3rd) dose for <u>immunocompromised</u> , at least 28 days after 2nd Pfizer dose	1		
Pfizer/	Primary 2 doses (21 days apart)		1	1
Comirnaty (12+)	Additional (3rd) dose for <u>immunocompromised</u> , at least 28 days after 2nd Pfizer dose		1	1
	Booster dose* of Pfizer (or Moderna if age 18+), at least 5 months after 2nd dose of Pfizer		1	1
Moderna (18+)	Primary 2 doses (28 days apart)			1
	Additional (3rd) dose for <u>immunocompromised</u> , at least 28 days after 2nd Moderna dose			1
	Booster dose* of Moderna (half-dose) or Pfizer, at least 6 months after 2nd dose of Moderna			1
Johnson & Johnson (18+) Use of Pfizer or Moderna is preferred.*	Primary 1 dose			1
	Booster dose* of different brand (preferred) or J&J, at least 2 months after primary J&J dose			1